



permobil

Case report references – John, Progeo and ROHO Client

References

1. Brienza, D., Vallely, J., Karg, P., Akins, J., & Gefen, A. (2018). An MRI investigation of the effects of user anatomy and wheelchair cushion type on tissue deformation. *Journal of Tissue Viability*, 27(1), 42–53. <https://doi.org/10.1016/j.jtv.2017.04.001>
2. Call, E., Hetzel, T., McLean, C., Burton, J. N., & Oberg, C. (2017). Off loading wheelchair cushion provides best case reduction in tissue deformation as indicated by MRI. *Journal of Tissue Viability*, 26(3), 172–179.
3. Crane, B., Winingar, M., & Call, E. (2016). Orthotic-Style Off-Loading Wheelchair Seat Cushion Reduces Interface Pressure Under Ischial Tuberosities and Sacrococcygeal Regions. *Archives of Physical Medicine and Rehabilitation*, 97(11), 1872–1879. <https://doi.org/10.1016/j.apmr.2016.04.004> <https://doi.org/10.1016/j.jtv.2017.05.002>
4. Hosseini, S. M., Oyster, M. L., Kirby, R. L., Harrington, A. L., & Boninger, M. L. (2012). Manual wheelchair skills capacity predicts quality of life and community integration in persons with spinal cord injury. *Archives of physical medicine and rehabilitation*, 93(12), 2237-2243.
5. Koo, T. K., Mak, A. F., & Lee, Y. L. (1995). Evaluation of an active seating system for pressure relief. *Assist Technol*, 7(2), 119-128. doi:10.1080/10400435.1995.10132260
6. May, L. A., Butt, C., Kolbinson, K., Minor, L., & Tulloch, K. (2004). Wheelchair back-support options: Functional outcomes for persons with recent spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 85(7), 1146–1150. <https://doi.org/10.1016/j.apmr.2003.08.105>
7. Requejo, P. S., Furumasu, J., & Mulroy, S. J. (2015). Evidence-based strategies for preserving mobility for elderly and aging manual wheelchair users. *Topics in geriatric rehabilitation*, 31(1), 26.
8. Presperin Pedersen, J., Smith, C., Dahlin, M., Henry, M., Jones, J., McKenzie, K., Sevigny, M., & Yingling, L. (2020). Wheelchair backs that support the spinal curves: Assessing postural and functional changes. *The Journal of Spinal Cord Medicine*, 1–10. <https://doi.org/10.1080/107>
9. Sawatzky, B., DiGiovine, C., Berner, T., Roesler, T., & Katte, L. (2015). The need for updated clinical practice guidelines for preservation of upper extremities in manual wheelchair users: a position paper. *American journal of physical medicine & rehabilitation*, 94(4), 313-324.
10. Sprigle, S., Wootten, M., Sawacha, Z., & Thielman, G. (2003). Relationships among cushion type, backrest height, seated posture, and reach of wheelchair users with spinal cord injury. *J Spinal Cord Med*, 26(3), 236-243. doi:10.1080/10790268.2003.11753690
11. Thyberg, M., Gerdle, B., Samuelsson, K., & Larsson, H. (2001). Wheelchair seating intervention. Results from a client-centred approach. *Disability and Rehabilitation*, 23(15), 677-682.
12. Worobey, L. A., Bernstein, J., Ott, J., Berner, T., Black, J., Cabarle, M., ... & Betz, K. (2023). RESNA position on the application of ultralight manual wheelchairs. *Assistive Technology*, 1-18.